

Literatur

- BERRY, P. Y. (1964): The Breeding Patterns of Seven Species of Singapore Anura. – *Journal of Animal Ecology* 33 (2): 227–243.
- BOLKAY, S. J. (1915): Beiträge zur Osteologie einiger exotischer Raniden. – *Anatomischer Anzeiger* 48: 172–183.
- DASH, M. C. & J. K. MAHANTA (1993): Quantitative analysis of the community structure of tropical amphibian assemblages and its significance to conservation. – *J. Biosci.* 18 (19): 121–139.
- DUBOIS, A., & A. ÖHLER (2000): Systematics of *Fejervarya limnocharis* (GRAVENHORST, 1829) (Amphibia, Anura, Ranidae) and related species. 1. Nomenclatural status and type-specimens of the nominal species *Rana limnocharis* GRAVENHORST, 1829. – *Alytes* 18: 15–50.
- FARUK A., D. BELABUT, N. AHMAD, R. J. KNELL & T. W. GARNER (2013): Effects of oil-palm plantations on diversity of tropical anurans. – *Conserv Biol.* 27: 615–24.
- FROST, D. R. (2018): Amphibian Species of the World: an Online Reference. Version 6.0. Online unter <http://research.amnh.org/herpetology/amphibia/index.html>. – American Museum of Natural History, New York, USA. Download: 20.08.2018
- GRAVENHORST, J. L. C. (1829): Deliciae Musei Zoologici Vratislaviensis. Fasciculus primus. Chelonios et Batrachia. – Leipzig: Leopold Voss.
- HAILEY, A., N. SOOKOO, A. MOHAMMED & A. KHAN (2006): Factors affecting tadpole growth: Development of a rearing system for the Neotropical leptodactylid *Physalaemus pustulosus* for ecotoxicological studies. – *Appl. Herpetol.* 3: 111–128.
- , –, E. HERNANDEZ & D. RAMOO (2007): The influence of density and ration level on cultured *Physalaemus pustulosus* tadpoles, and the mitigation of a crowding effect by soil substrate. – *Applied Herpetology* 4: 261–277.
- ISKANDAR, D. T. (1998): The Amphibians of Java and Bali. – Bogor, Indonesia: Research and Development Centre for Biology – LIPI and GEF Biodiversity Collections Project.
- JAAFAR, I. H. (1994): The life history, population and feeding biology of two paddy field frogs, *Rana cancrivora* GRAVENHOS and *R. limnocharis* BOIE, in Malaysia. – Doktorarbeit, Universität Pertanian Malaysia.
- A. ISMAIL & A.-R. KURAI (1999): Correlations of Reproductive Parameters of Two Tropical Frogs from Malaysia. – *Asiatic Herpetological Research*, 8: 48–52.
- KÖHLER, G., L. MOGK, K.P.P. KHAING & N.L. THAN (2019): The genera *Fejervarya* and *Minervarya* in Myanmar: Description of a new species, new country records, and taxonomic notes (Amphibia, Anura, Dicroglossidae). – *Vertebrate Zoology* 69 (2): 183–226
- KOTAKI, M., A. KURABAYASHI, M. MATSUI, M. KURAMOTO, T.H. DJONG & M. SUMIDA (2010): Molecular Phylogeny of the Diversified Frogs of Genus *Fejervarya* (Anura: Dicroglossidae). – *Zoological Science*, 27: 386–395.
- KUSRINI, M. D. (2005): Edible frog harvesting in Indonesia: Evaluating its impact and ecological context. – Doktorarbeit, James Cook University.
- KUNZ, K. (2015): Schaumschläger und Topmodel. Spannendes rund um den Túngara-Frosch. Teil 2. – *TERRARIA/elaphe* 56: 56–69.
- MIN, P. Y. (2007): Comparative aspects of metamorphosis in *Fejervarya limnocharis* and *Fejervarya cancrivora* (Amphibia: Anura). – Bachelor-Arbeit, Universiti Malaysia Sarawak.
- OTHMAN, M. S., W. KHONSUE, J. KITANA, K. THIRAKHUP, M. G. ROBSON & N. KITANA (2011): Reproductive Mode of *Fejervarya limnocharis* (Anura: Ranidae) Caught from Mae Sot, Thailand Based on Its Gonadosomatic Indices. – *Asian Herpetol Res.* 2(1): 41–45.
- PHAM, C. T., H. T. AN, S. HERBST, M. BONKOWSKI, T. ZIEGLER & T. Q. NGUYEN (2017): First report on the amphibian fauna of Ha Lang karst forest, Cao Bang Province, Vietnam. – *Bonn zoological Bulletin* 66 (1): 37–53.
- PREMO, D. B. (1985): The reproductive ecology of a ranid frog community in pond habitats of West Java, Indonesia. – Doktorarbeit, Michigan State University.
- TARKHNISHVILI, D. (1994): Amphibian communities of the southern Viet Nam: preliminary data. – *Journal of Bengal Natural History Society, New Series*, 1 (13): 3–62.
- TODA, M., M. MATSUI, M. NISHIDA & H. OTA (1998a): Genetic divergence among Southeast and East Asian populations of *Rana limnocharis* (Amphibia: Anura), with special reference to sympatric cryptic species in Java. – *Zool. Ser.*, 15: 607–613.
- , M. NISHIDA, M. MATSUI, W. G. FU & H. OTA (1997): Allozyme variation among East Asian populations of the Indian rice frog, *Rana limnocharis* (Amphibia: Anura). – *Biochem. Syst. Ecol.*, 25: 143–159.
- , –, K. Y. LUE & H. OTA (1998b): Genetic variation in the Indian rice frog, *Rana limnocharis* (Amphibia: Anura), in Taiwan, as revealed by allozyme data. – *Herpetologica*, 54 (1): 73–82.
- ZIEGLER, T. (2002): Die Amphibien und Reptilien eines Tieflandfeuchtwald-Schutzgebietes in Vietnam. – *Natur und Tier - Verlag, Münster*.
- & S. WEITKUS (1999): Zur Herpetofauna eines nordvietnamesischen Küstenbiotops: Einnischung und Mageninhaltsuntersuchungen. Teil 2. – *herpetofauna*, 21 (123): 25–33.